

Research Article



CrossMark

Open Access

2002년 금강, 만경-동진강 하천수 중 잔류농약의 연간 검출 양상

김찬섭*, 이희동, 임양빈, 손경애

가

Temporal Patterns of Pesticide Residues in the Keum, Mangyung and Dongjin Rivers in 2002

Chan-sub Kim*, Hee-Dong Lee, Yang-Bin Ihm and Kyeong-Ae Son (Agromaterial Assessment Division, Department of Agro-food Safety & Crop Protection, National Institute of Agricultural Sciences, Rural Development Administration, Wanju 55365, Korea)

Received: 17 November 2017/ Revised: 24 November 2017/ Accepted: 26 November 2017

Copyright © 2017 The Korean Society of Environmental Agriculture

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Chan-sub Kim

<http://orcid.org/0000-0003-2157-7311>

Abstract

BACKGROUND: To evaluate residues of environmentally concerned pesticides in water system, this monitoring was conducted over three rivers. The residual characteristics and discharging condition of these residues on water system was investigated.

METHODS AND RESULTS: Total twenty nine sampling sites were selected through main streams and branch streams of Keum, Mangyung and Dongjin rivers, and the water samples from them were regularly collected one month interval, especially biweekly from May to August in 2002. Of the pesticides monitored, six fungicides which include hexaconazole, isoprothiolane and iprobenfos were detected with frequencies of 0.3-50.9% and in their residue level of 0.1-4.7 µg/L. Sixteen insecticides which include nine organophosphoruses, three carbamates, endosulfan, cypermethrin, buprofezin and fipronil were detected with frequencies of 0.3-32.5% and in their residue level of 0.01-2.8 µg/L. Nine herbicides which include alachlor molinate, anilofos, butachlor, dimepiperate, metolachlor, oxadiazon, pretilachlor and thiobencarb were detected with frequencies of 0.8-22.9% and in their residue level of 0.01-9.07 µg/L.

CONCLUSION: Detection frequencies and residue levels of insecticides and herbicides were the highest in waters sampled in May and June. Almost pesticides detected were for the paddy rice and their residue levels were very low to compare with standard values.

Key words: Monitoring, Paddy rice, Pesticide residues, River water

서 론

가 , 가 (Masiá *et al.*, 2015). 가 (Ccanccapa, *et al.*, 2016b; Konstantinou *et al.*, 2006).

가 . 가 가 .

*Corresponding author: Chan-Sub Kim
Phone: +82-63-238-3358; Fax: +82-63-238-3839;
E-mail: chskim@korea.kr

가 .

2002
3-11

가

(Lee *et al.*, 2011; Papadakis *et al.*, 2015a and 2015b; Ccanccapa, *et al.*, 2016b),

가

(Konstantinou *et al.*, 2006; Masiá *et al.*, 2013 and 2015; Ccanccapa, *et al.*, 2016a).

1970 (Park & Park, 1980; Park & Ma, 1982; Choi *et al.*, 2011), (Park & Hwang, 1982; Lee *et al.*, 1985), 가 가

1983 and 1984; You & Park, 1984, Park *et al.*, 1996), 가

(Lee *et al.*, 1976; Yu *et al.*, 2002) 가

(Lee *et al.*, 1976; Park & Hwang, 1982; Suh *et al.*, 1986) (Lee *et al.*, 1976; Park & Hwang, 1982; Choi *et al.*, 2011)

재료 및 방법

시료채취

(Lee *et al.*, 1983 and 1984), 16, 7, 6

29, 1

2-3 5~8 2

가 6-8 가

가 1

가 (




Fig. 1. Map of sampling sites for monitoring of pesticide residues in Keum rivers; asterik is station of main stream, circle station of branch stream.

Table 1. GLC conditions for pesticide residue analysis

Gas chromatograph:	HP 5890 series II plus with 7673 auto-sampler		
Detector:	ECD (Electron capture detector) NPD (Nitrogen phosphorus detector)		
Capillary Column:	Guard column, 10 m × 0.25 mm ID and DB-5, 30 m × 0.25 mm ID (film thickness 0.25 µm)		
Temperature:	Detector	ECD	300°C
		NPD	270°C
	Injection port		230°C
Flow:	Column oven	60°C (2min) — 20°C/min → 120°C — 5°C/min → 270°C (15 min)	
	Carrier	Helium	30cm/sec
	Fuel (NPD)	Hydrogen	3.5 mL/min
Sampling mode:		Air	100 mL/min
	Make-up ECD	Nitrogen	60 mL/min
	NPD	Nitrogen	30 mL/min
Sample volume:	ECD	1 µL	
	NPD	2 µL	

Table 2. HPLC conditions for carbendazim, imidacloprid and acetamiprid residue analysis

HPLC:	HP 1100 series with auto-sampler
Column:	C18 column (HP Zorbax XDB18, 4.6 mm×25 cm)
Detector:	Diode-array detector, 246, 278, 285 nm
Mobile phase:	CH ₃ CN/methanol/0.02 M potassium phosphate buffer (pH 7.0) (17/8/75)
Flow rate:	1 mL/min
Injection volume:	20 µL

Table 3. Summary of some fungicide residues in river waters

Pesticide	No. of samples detected	Main season detected (month)	Concentration (µg/L)				Standard (µg/L)
			in positive samples			Average of total samples	
			Range	Average	Median		
Carbendazim	7	7~8	0.5~7.7	1.6	0.6	0.031	200 ^{a)}
Vinclozolin	1	8	0.2	0.2	0.2	0.0006	-
Hexaconazole	30	7~10	0.1~2.9	0.6	0.3	0.045	-
Iprobenfos	108	6~8	0.1~4.3	0.3	0.2	0.096	5 ^{b)}
Isoprothiolane	191	5~10	0.3~4.7	0.9	0.6	0.475	40 ^{b)}
Edifenphos	10	7~8	0.1~0.2	0.1	0.1	0.003	6 ^{c)}

^{a)}Australia, drinking water standards, ^{b)}Japan, drinking water standards ^{c)}Japan, guideline value for environmental assessment.

1 g

acetonitrile 5 mL

결과 및 고찰

살균제

3

11

기기분석

Table 3

Fig. 3

WWF

GLC

가

carbendazim

vinclozolin,

Table 1

(Kim *et al.*, 2010), GC/MS

hexaconazole

Finnigan

GCQ

GLC

iprobe

isoprothiolane, edifenphos

, HPLC

Table 2

Carbendazim

benomyl




Fig. 3. Seasonal changes of some fungicide residue levels in river waters.

thiophanate-methyl . 3 MT
 benomyl carbendazim 1 (, 20003),
 , thiophanate-methyl 10
 (Pesticide manual, 2012) benomyl 가 (, 2002). 1992
 carbendazim 376 (0.06 µg/L)
 1.9% 7 0.5 µg/L (, 1992).
 5 800 MT(, 20003) Hexaconazole 8 9 29
 1.6 µg/L , 11 1 µg/L 20
 7.7 µg/L 0.5-0.8 µg/L 2/3 (hexaconazole
 1998 56 , , 1998)
 (4 8 ,
 1 0.8 µg/L 0.9 µg/L
 g/L 8 1 0.6 µg/L isoprothiolane iprobenfos, edifenfos
)(, 1998). 7~8
 200 µg/L 4.7, 4.3 0.2 µg/L ,
 1/7000 Catalonia 191, 108 10 2001
 2011 0.0108-0.6974 µg/L 21% iprobenfos 1,230 MT, isoprothiolane 550
 (Masiá, A *et al.*, 2015), MT edifenfos 110 MT (, 2003),
 2010-2013 0-25% isoprothiolane
 0.0020-0.0063 µg/L (Ccanccapa *et al.*, 2016a), 가 (Pesticide
 Gadalquivir 2011 0.0006-0.0114 manual, 2012). Isoprothiolane 4.7 µg/L
 µg/L 17% (Masiá, A *et al.*, 2013), 0.475 µg/L Lee (1984) 1983
 Ebro 2011 0.00004-0.0116 µg/L 17% 12 2
 (Ccanccapa *et al.*, 2016b) 6 8 0.96 µg/L
 0.281 µg/L, (,
 Vinclozolin 376 1992) 6, 8 0.2-2.8 µg/L
 8 22 0.2 µg/L (, 1998) 6, 8
 . Vinclozolin 가 5 0.3-11.0 µg/L . 가

Table 4. Summary of some insecticide residues in river waters

Pesticide	No. of samples detected	Main period detected (month)	Concentration (μg/L)			Average of total samples	Standard (μg/L)
			in positive samples				
			Range	Average	Median		
Endosulfan	122	6~10	0.02~1.06	0.13	0.09	0.043	40 ^{a)}
Cypermethrin	1	8	1.7	1.7	1.7	0.0045	-
Fenoxycarb	2	8	0.5~0.7	0.6	0.6	0.0032	-
Buprofezin	9	8	0.02~0.09	0.05	0.06	0.0013	40 ^{b)}
Carbofuran	53	5~8	0.1~2.8	0.4	0.3	0.056	40 ^{c)}
Diazinon	96	4~8	0.01~0.33	0.05	0.03	0.013	20 ^{d)}
EPN	7	8	0.1~0.2	0.2	0.1	0.0024	6 ^{e)}
Ethoprophos	16	5~6	0.1~0.5	0.3	0.2	0.011	-
Fenobucarb	10	7~8	0.1~0.5	0.2	0.2	0.0053	20 ^{e)}

^{a)}Australia, drinking water standards, ^{b)}Japan, guideline value for environmental assessment, ^{c)}US EPA, national primary drinking water standards, ^{d)}Korea, drinking water standards, ^{e)}Japan, drinking water standards.




Fig. 4. Seasonal changes of some insecticide residue levels in river waters.

1982 2012 isoprothiolane 0.38 µg/L), (, 1992)
320-740 MT 가 6, 8 0.05-2.21 µg/L
(, 1987-2003). (, 1998) 6, 8
iprobefos 4.3 µg/L 0.4-4.2 µg/L . 1980
5 µg/L 가 가 2000
800-1560 MT(, 1987-2003)
가 . 1982 .
<0.05-1.53 µg/L 0.02 Edifenfos 7, 8 0.1-0.2 µg/L
µg/L (4), 0.41 µg/L (8) (Lee *et al*, Lee (1984)
8 0.222 µg/L 0.086 µg/L , edifenfos
(Lee *et al*, 1984). 1990 1990 250-460 MT가
(1991,) iprobefos (, 1987-2003).
(<0.05-0.73 µg/L 4 0.06 µg/L, 8

Table 5. Summary of some herbicide residues in river waters

Pesticide	No. of samples detected	Main period detected (month)	Concentration (µg/L)				Standard (µg/L)
			in positive samples			Average of total samples	
			Range	Average	Median		
Alachlor	29	4~5, 7	0.1~1.9	0.5	0.5	0.042	2 ^{a)}
Molinate	86	5~7	0.01~9.07	1.66	0.56	0.381	7 ^{b)}
Anilofos	5	6	0.1~0.2	0.1	0.1	0.0016	-
Butachlor	18	5~6	0.1~0.9	0.4	0.4	0.021	30 ^{c)}
Dimepiperate	3	6	0.2~0.5	0.3	0.3	0.0027	-
Metolachlor	9	4~5	0.4~0.9	0.6	0.5	0.014	5 ^{b)}
Oxadiazon	10	5~6	0.2~0.5	0.3	0.3	0.009	-
Pretilachlor	5	5~6	0.4~1.1	0.6	0.5	0.008	-
Thiobencarb	24	5~6	0.1~3.6	0.8	0.4	0.049	20 ^{d)}

^{a)}US EPA, national primary drinking water standards, ^{b)}WHO, guideline value recommended, ^{c)}Australia, drinking water standards, ^{d)}Japan, drinking water standards.




Fig. 5. Seasonal changes of some herbicide residue levels in river waters.

-0.0044 µg/L 14% 2011 Lee (1984) 8-12
(Masiá, A *et al.*, 2015), 4 0.23-0.95 µg/L, (
2010-2013 0-15% 1992) 6 1 , 8 16 0.5-1.2 µg/L,
<0.00001-0.013 µg/L (Ccanccapa *et al.*, (, 1998) 1998 8
2016a), Gadalquivir 2010 0.3 µg/L fenobucarb
0.0018-0.0093 µg/L 23% 2011
(Masiá, A *et al.*, 2013),
Ebro 2010 0.0023-0.0083 µg/L 22% 제초제
2011 (Ccanccapa
et al., 2016b) . Alachlor , , pendimethalin
Fenobucarb 0.1~0.5 µg/L 7-8 가 , , , , ,
10 , (, 2003),

5	400-500M/T	가	(5	7	.
2003).		376	7.7%			6
29	0.1-1.9 µg/L				molinate가	,
	0.5 µg/L	.	(7		
	, 1998)	2.6%	3			(
		0.37 µg/L			, 1998)	1998 6
				0.3-40 µg/L	73.7%	,
4	5			(Molinat	, 2001)
, 7				2000 6-8		6
		2010-2013		53.4%, 7	20.7%	39.4 µg/L
	2013 1	<0.00001 µg/L			. Konstantinou	(2006) 1988-2000
(Ccanccapa <i>et al.</i> , 2016a)					1990-2000	4
Konstantinou (2006) 1988-2000					-	molinate
2000 5					0.036-0.900 µg/L	0.038-4.8000 µg/L
-	alachlor	0.031- 9.300 µ			376 23%	86
g/L 0.014-0.206 µg/L		. Alachlor		0.01-9.1 µg/L		
		1-66 µg/L		1.7 µg/L		가
0.001%	(Kim <i>et al.</i> , 2006b),					
	0.43-0.81 µg/L	(EPA			26
가 , 1998)		alachlor		2-5		
	metolachlor	4~5	7			
0.4-0.9 µg/L					가	
9	alachlor 1/3				1.4-1.9 µg/L	
Catalonia	2010-2011	0.0016-0.0130				
µg/L 14%	(Masiá, A <i>et al.</i> , 2015),				가 2.2-8.3 µg/L	
	2010-2013	0-23%				4
	<0.00001-0.0022 µg/L	(Ccanccapa		WHO	7 µg/L	
<i>et al.</i> , 2016a),	Ebro 2010					
	2011 0.0011-0.0486 µg/L					
29%	(Ccanccapa <i>et al.</i> , 2016b)			Thiobencarb	butachlor	
. Konstantinou (2006) 1988-2000					5	6
11990-2000	5					, , ,
-	metolachlor				6	
0.060-1.120 µg/L	0.004-3.000 µg/L					(
Metolachlor		0.09-			, 1998)	butachlor thiobencarb,
1.01 µg/L	0.20-1.86%	(Kim <i>et al.</i> ,		oxadiazon 6	28.9% 18.4%, 26.3%	
2006a),		2.4%		0.4-2.9 µg/L	0.3-1.5 µg/L, 0.2-0.5 µg/L	
	(Gaynor <i>et al.</i> , 1995),					
	가	10 µg/L	(butachlor 7	0.1-0.9 µg/L,
EPA	가 , 1995).			thiobencarb 5	0.2-0.5 µg/L, oxadiazon	
		alachlor		4	0.2-0.5 µg/L	pretilachlor 3
가		4~51 µg/L		0.5-1.1 µg/L		
4	alachlor					
		가				
	alachlor가					가
가						
Molinate						isoprothiolane

결 론

endosulfan

가 가 , molinate

요 약

가

8,90

11 1 , 5-8

2 carbendazim

hexaconazole 6 0.3-

50.9% 0.1-4.7 µg/L

isoprothiolane iprobenfos

endosulfan 16

0.3-32.5% 0.01-2.8 µ

g/L endosulfan 28

isoprothiolane

iprobenfos, endosulfan

가 가

alachlor 9

0.8-22.9% 0.01- 9.07 µg/L

. Molinate 4 WHO

7 µg/L

Notes

The author declare no conflict of interest.

Acknowledgement

This study was carried out with the support of Research Program for Agricultural Science & Technology Development ("Monitoring for endocrine disrupting pesticide residues in river waters in 2002" and Project No. PJ012542012017), National Institute of Agricultural Sciences, Rural Development Administration, Republic of Korea.

References

- Ccanccapa, A., Masiá, A., Andreu, V., & Picó, Y. (2016a). Spatio-temporal patterns of pesticide residues in the Turia and Júcar Rivers (Spain). *Science of the Total Environment*, 540, 200-210.
- Ccanccapa, A., Masiá, A., Navarro-Ortega, A., Picó, Y., & Barceló, D. (2016b). Pesticides in the Ebro River basin: Occurrence and risk assessment. *Environmental Pollution*, 211, 414-424.
- Choi, J. Y., Lee, S. G., Bang, J. H., Yang, D. B., Hong, G. H., & Shin, K. H. (2011). On the distribution of PCBs and organochlorine pesticides in fish and sediment of the Asan bay. *Ocean and Polar Research*, 33(1), 45-53.
- Gaynor, J. D., MacTavish, D.C., & Findlay, W. I. (1995). Atrazine and metolachlor loss in surface and subsurface runoff from three tillage treatments in corn, *Journal of Environmental Quality*, 24, 246-256.
- Kim, C. S., Lee, B. M., Park, B. J., Jung, P. K., Choi, J. H., & Ryu, G. H. (2006a). Runoff of diazinon and metolachlor by rainfall simulation and from soybean-grown field lysimeter. *Korean Journal of Pesticide Science*, 10(4), 279-288.
- Kim, C. S., Lee, H. D., Oh, B. Y., & Lee, Y. D. (2006b). Runoff and erosion of alachlor, ethalfuralin, ethoprophos and pendimethalin from soybean field lysimeter. *Korean Journal of Environmental Agriculture*, 25(4), 297-305.
- Kim, C. S., Lee, H. D., Ihm, Y. B., & Im, G. J. (2007). Runoff of endosulfan by rainfall simulation and from soybean-grown field lysimeter. *Korean Journal of Environmental Agriculture*, 26(4), 343-350.
- Kim, C. S., Lee, B. M., Park, K. H., Park, B. J., Park, J. E., & Lee, Y. D. (2010). Simultaneous determination of pesticide residues in soils by dichloromethane partition - adsorption chromatography - GC-ECD/NPD analytical methods. *Korean Journal of Pesticide Science*, 14(4), 361-370.
- Konstantinou, I. K., Hela, D. G., & Albanis, T. A. (2006). The status of pesticide pollution in surface waters (rivers and lakes) of Greece. Part I. Review on occurrence and levels. *Environmental Pollution*, 141(3), 555-570.
- Lee, S. R., Kang, S. Y., Park, C. K., Lee, J. H., & Rho, C. S. (1976). A survey on the residues of organochlorine pesticide in water, mud and clam samples from the Kwangyang bay, Korea. *Journal of Korean Agricultural Chemical Society*, 19(2), 112-118.
- Lee, H. K., Lee, Y. D., Park, Y. S., & Shin, Y. H. (1983). A survey for pesticide residues in major Rivers of Korea. *Korean Journal of Environmental Agriculture*,

- 2(2), 83-89.
- Lee, H. K., Lee, Y. D., & Park, Y. S. (1984). Evaluation of pesticide residues of river waters in 1983. Research Report of ORD (S.P.M.U), 26(1), 46-53.
- Lee, Y. H., Hwang, E. C., & Park, C. K. (1985). Evaluation of polychlorinated biphenyls (PCBs) and organochlorine insecticide residues in irrigation waters in the periphery of Suwon. Korean Journal of Environmental Agriculture, 4(2), 95-101.
- Lee, J. H., Park, B. J., Park, S. W., Kim, W. I., Hong, S. M., Im, G. J., & Hong, M. G. (2011). Ecological risk assessment of pesticide residues in agricultural lake: Risk quotients and probabilistic approach. Korean Journal of Environmental Agriculture, 30(3), 316-322.
- Masiá, A., Campo, J., Vázquez-Roig, P., Blasco, C., & Picó, Y. (2013). Screening of currently used pesticides in water, sediments and biota of the Guadalquivir River Basin (Spain). Journal of Hazardous Materials, 263(Part-1), 95-104.
- Masiá, A., Campo, J., Navarro-Ortega, A., Barceló, D., & Picó, Y. (2015). Pesticide monitoring in the basin of Llobregat River (Catalonia, Spain) and comparison with historical data. Science of the Total Environment, 503-504, 58-68.
- Papadakis, E. N., Tsaoulas, A., Kotopoulou, A., Kintzikoglou, K., Vryzas, Z., & Papadopoulou-Mourkidou, E. (2015a). Pesticides in the surface waters of Lake Vistonis basin, Greece: Occurrence and environmental risk assessment. Science of the Total Environment, 536, 793-802.
- Papadakis, E. N., Vryzas, Z., Kotopoulou, A., Kintzikoglou, K., Makris, K. C., & Papadopoulou-Mourkidou, E. (2015b). A pesticide monitoring survey in rivers and lakes of northern Greece and its human and ecotoxicological risk assessment. Ecotoxicology and Environmental Safety, 116, 1-9.
- Park, C. K., & Park, N. D. (1980). Analysis of organochlorine pesticide residues in the presence of polychlorinated biphenyls (PCBs), Part II. Analysis of river sediments and cultivating soils in the peripheries of several industrial estates. Journal of Korean Agricultural Chemical Society, 29(1), 58-63.
- Park, C. K., & Ma, Y. S. (1982). Organochlorine pesticide residues in agricultural soils - 1981. Korean Journal of Environmental Agriculture, 1(1), 1-13.
- Park, C. K., & Hwang, E. C. (1982). Evaluation of polychlorinated biphenyls and organochlorine insecticide residues in water, sediment and Crusian carps in Soho lake. Korean Journal of Environmental Agriculture, 1(2), 105-115.
- Park, Y. K., Lee, C. H., Lee, S. H., & Kim, J. W. (1996). Characteristics of pesticide discharge in the Nakdong river basin: Evaluation of pesticide discharge from farmland and golf links. Journal of Korean Society of Environmental Engineers, 18(5), 627-636.
- Suh, Y. T., Im, G. J., & Shim, J. H. (1986). Evaluation of organochlorine pesticide residues in the mud flat. Korean Journal of Environmental Agriculture, 5(2), 113-118.
- You, S. J., & Park, C. K. (1984). On the organochlorine pesticide residues in downstream area of Nakdong river. Journal of Korean Agricultural Chemical Society, 27(3), 187-197.
- Yu, J., Yang, D. B., Kim, K. T., & Lee, K. W. (2002). Distribution of organophosphorus pesticides in Asan and Kyeonggi bay, Korea. Journal of Korean Society of Marine Environmental Engineering, 5(1), 38-50.