

Research Article



CrossMark

Open Access

야관문 에탄올 추출물의 항산화 및 Metalloproteinase 저해 활성

신용하, 송창길*

Antioxidant and Metalloproteinase Inhibitory Activities of Ethanol Extracts from *Lespedeza cuneata* G. Don

Yong Ha Shin and Chang-Khil Song* (Department of Agriculture, Graduate School of Jeju National University, Jeju 63243, Korea)

Received: 23 October 2017/ Revised: 30 October 2017/ Accepted: 1 December 2017

Copyright © 2017 The Korean Society of Environmental Agriculture

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Yong Ha Shin

<http://orcid.org/0000-0002-5896-0560>

Chang-Khil Song

<http://orcid.org/0000-0002-5394-8378>

Abstract

BACKGROUND: *Lespedeza cuneata* G. Don is a well-known medicinal plant. In this study, the biological activities of *L. cuneata* extracts were investigated.

METHODS AND RESULTS: *L. cuneata* shoot was extracted with 30% ethanol and further fractionated with organic solvents. Total phenolic and flavonoid content, antioxidant activity, and matrix metalloproteinase inhibition effect of the extract and fractions were measured. Among the tested extract and fractions, the highest contents of total phenolic and flavonoids were found in ethyl acetate fraction (117.8 mg GAE/g and 35.9 mg QE/g, respectively). Ethyl acetate fraction showed the highest DPPH and ABTS radical scavenging activity, and the antioxidant activity of the other fractions followed the order n-hexane fraction>ethanol extract>methyl chloroform>n-butanol fraction. Inhibitory effect on the expression of matrix metalloproteinases (MMP1 and MMP3) was highest in the fraction of ethyl acetate, and n-butanol fraction also significantly inhibited the expression of MMP3. Antioxidant activities of *L. cuneata* extracts were significantly positively related to their phenolic and flavonoid content.

CONCLUSION: Ethyl acetate fraction of *L. cuneata* ethanol extract showed potent antioxidant and matrix metalloproteinases inhibitory activities. Those activities might be related to the high total phenolic and flavonoid content of the extract.

Key words: Antioxidant, Flavonoid, *Lespedeza cuneata*, Matrix metalloproteinase, Phenol

서론

(*Lespedeza cuneata* G. Don)

가

al, 2014).

가

1991).

(collagen)

(elastin)

. Collagen matrix metalloproteinase-1 (MMP1) elastin

(Jung et

가

(Stich,

*Corresponding author: Chang-Khil Song
Phone: +82-64-710-7585; Fax: +82-64-725-2351;
E-mail: sck5622@jejunu.ac.kr

matrix metalloproteinase-3 (MMP3) 가 (Rhie *et al.*, 2001; Tsuji *et al.*, 2001). MMPs 가 collagen elastin 가 MMPs (Stich, 1991). flavonoid phenol quercetin kaempferol (Numata *et al.*, 1980). tyrosinase ethyl acetate 가 (Cha *et al.*, 2010; Lee *et al.*, 2011). elastase 가 (Kim and Kim, 2014). collagenase elastinase human dermal fibroblast (HDF)

gallic acid equivalent(GAE) . Kim (2011) 200 µL 10% aluminum nitrate 100 µL 1 M potassium acetate 100 µL 4.6 mL 80% 가 417 nm 3 quercetin quercetin equivalent(QE) **항산화 활성 측정** DPPH ABTS radical DPPH radical Kim (2011) 20 µL 0.15 mM 1,1-diphenyl-2-picrylhydrazyl(DPPH) 180 µL 가 96-well plate 30 multi-micro plate reader (Spectra Max i3X, CA, USA) 517 nm 3 DPPH radical , DPPH radical 50% DRC50

재료 및 방법

시료 및 추출물의 제조

500 g 30% (v/v) 1:15 24 n-hexane, methyl chloroform, ethyl acetate, n-butanol, aqueous 3

$$\text{Inhibition rate (\%)} = [1 - (A - B) / C] \times 100$$

A: absorbance of sample

B: absorbance of blank

C: absorbance of control

ABTS radical Ko (2017) 7.4 mM 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS) 2.6 mM potassium persulphate ABTS

총 페놀 및 플라보노이드 함량 분석

Folin-Ciocalteu (Kim *et al.*, 2011). 30% 100 µL 500 µL Folin-Ciocalteu 50 µL 5 20% sodium carbonate 300 µL 20 725 nm 3

ABTS 가 734 nm 가 1.0 20 µL multi-microplate reader 734 nm 3 , ABTS radical 50% ARC50

MMP1과 MMP3 저해 활성 측정

MMPs Cho (2016) Human dermal

fibroblast (HDF) 1×10^5 cells/well
 6-well plate tumor necrosis factor alpha
 (TNF- α) 10 ng/mL MMPs
 48 phosphate buffered
 saline
 protein extraction solution 가 1.5
 mL tube 99°C 10 가
 -70°C bicinchoninic acid protein
 assay kit 10% SDS
 polyacrylamide gel nitrocellulose membrane
 membrane 5% skim milk/TBST
 (tris-buffered saline and 0.1% Tween 20) 1
 blocking 1 antibody membrane 가
 가 TBST 5 5 가
 1 antibody peroxidase 2 가
 antibody 1
 Membrane EZ-Western lumi pico reagent 가
 ChemiDoc image analyzer(Bio-Rad)
 ImageJ program 가
 (Kim *et al.*, 2011; Senguttuvan *et al.*, 2014),

통계 처리
 test ,
 가 .
 Duncan's multiple range
 Peason's correlation coefficient

결과 및 고찰
 페놀, 플라보노이드 함량
 Table 1
 가 , 30%
 가 ,
 가 ,
 가 , 30%

Table 1. Total phenol and flavonoid contents in 30% ethanol extract and soluble fractions of *Lespedeza cuneata* G. Don

Extract and Fractions	Total phenol (mg GAE/g) ^{a)}	Total flavonoid (mg QE/g) ^{b)}
30% Ethanol extract	36.21±0.24 e ^{c)}	5.13±0.54 e
<i>n</i> -Hexane fraction	79.74±0.78 b	6.36±1.20 d
Methyl chloroform fraction	39.06±1.10 d	7.93±0.43 c
Ethylacetate fraction	117.77±2.86 a	35.89±0.55 a
<i>n</i> -Butanol fraction	45.40±0.80 c	14.71±0.23 b
Aqueous fraction	13.31±0.57 f	Not detected

^{a)}GAE: gallic acid equivalent

^{b)}QE: quercetin equivalent

^{c)}Each value represents mean±SD (n=3), and same letters within a row indicate no significant difference at $p < 0.05$.

Table 2. Radical scavenging activity of 30% ethanol extract and soluble fractions of *Lespedeza cuneata* G. Don

Extract and Fractions	DRC ₅₀ (μ g/mL) ^{a)}	ARC ₅₀ (μ g/mL) ^{b)}
30% Ethanol extract	178.94±5.46 c ^{d)}	59.60±0.80 b
<i>n</i> -Hexane fraction	124.11±2.18 de	51.13±0.26 c
Methyl chloroform fraction	190.49±3.77 c	58.72±1.22 b
Ethylacetate fraction	91.63±1.96 e	26.82±0.61 d
<i>n</i> -Butanol fraction	157.38±4.17 cd	60.21±0.87 b
Aqueous fraction	833.24±63.1 a	161.36±8.14 a
BHT ^{c)}	241.35±8.48 b	< 6.25

^{a)}Concentration required to scavenge DPPH radical by 50%

^{b)}Concentration required to scavenge ABTS radical by 50%

^{c)}Positive control (Dibutylated hydroxytoluene),

^{d)}Each value represents mean±SD (n=3), and same letters within a row indicate no significant difference at $p < 0.05$.

가 . 가 . ABTS radical radical

항산화 활성

hydroxy toluene (BHT) . DPPH radical butylated (Zheng and Wang, 2001; Wojdylo *et al.*, 2007).

ABTS radical 50% 30% DRC₅₀

ARC₅₀ Table 2 가 ,

30% DPPH radical 가 ,

DPPH radical collagenase (MMP1, MMP8, MMP13), gelatinase (MMP2, MMP9), stromelysin (MMP3, MMP10) 3 , collagenase collagen collagen

Stromelysin collagen

ABTS radical 가

DPPH radical 가 MMP1 MMP3




Fig. 1. Effect of ethanol extract and soluble fractions of *Lespedeza cuneata* G. Don on MMP1 and MMP3 protein expression. Cells were stimulated by TNF- α for 48 hour in the presence of ethanol extract and soluble fractions of *Lespedeza cuneata* G. Don. Ex: ethanol extract, Hex: *n*-hexane fraction, MC: methyl chloroform fraction, EA: ethyl acetate fraction, Bu: *n*-butanol fraction, Aq: aqueous fraction.

Table 3. Correlation between total phenol and flavonoid content and biological activities of the ethanol extract and soluble fractions of *Lespedeza cuneata* G. Don

	TPC ^{a)}	TFC ^{b)}	DRC ₅₀ ^{c)}	ARC ₅₀ ^{d)}	MMP1	MMP3
TPC	1	0.8479***	-0.6461**	-0.7427**	-0.7687**	-0.8505***
TFC		1	-0.5289*	-0.6435**	-0.8269***	-0.8946***
DRC ₅₀			1	0.9841***	0.5003*	0.7775***
ARC ₅₀				1	0.6359**	0.8411***
MMP1					1	0.6948**
MMP3						1

^{a)}Total phenolic content determined as gallic acid equivalent

^{b)}Total flavonoid content determined as quercetin equivalent

^{c)}Concentration required to scavenge DPPH radical by 50%

^{d)}Concentration required to scavenge ABTS radical by 50%

Fig. 1 . 30%
 MMP1 MMP3
 가 가 , TNF- α
 control cell (-TNF- α) MMP1
 MMP3가 ,
 , MMP3 , 30%
 ,
 . 30% MMP1
 MMP3 .
 MMP1 MMP3
 (Kang *et al.*, 2007; Jung *et al.*, 2014),

상관관계 분석

Table 3 .
 radical DRC50 ARC50
 , collagen elastin
 MMP1 MMP3

요약

30% DPPH
 ABTS radical matrix metalloproteinase
 MMP1 MMP3
 가
 MMP1, MMP3
 가
 MMPs 가 가

Notes

The author declare no conflict of interest.

References

Cha, J. Y., Yang, H. J., Jeong, J. J., Seo, W. S., Park, J. S., Ok, M., & Cho, Y. S. (2010). Tyrosinase inhibition activity and antioxidant capacity by fermented products of some medicinal plants. *Journal of Life Science*, 20(6), 940-947.

Cho, Y. M., Song, H. S., Jang, S. A., Park, D. W., Shin, Y. S., Jeong, Y. J., & Kang, S. C. (2016). Suppression of VCAM-1 expression in human aortic smooth muscle cells treated with ethanol extracts of *Cynanchum wilfordii* Radix, *Arctium lappa* L., and *Dioscorea opposita*. *Korean Journal of Plant Resources*, 29(5), 525-531.

Jung, H. K., Choi, M. O., Kim, B. J., Jo, S. K., & Jeong, Y. S. (2014). Improving the efficacy of *Lespedeza cuneata* ethanol extract on ultraviolet-induced photoaging. *Korean Journal of Food Preservation*, 21(2), 264-275.

Kang, K. S., Kim, I. D., Kwon, R. H., Heo, Y. Y., Oh, S. H., Kim, M. A., Jung, H. J., Kang, H. Y., & Ha, B. J. (2007). The evaluation of anti-wrinkle effects in oriental herb extract. *Journal of Life Science*, 17(8), 1147-1151.

Kim, D. I., & Kim, H. J. (2014). Regeneration effects of *Lespedeza cuneata* ethanol extract on experimental open wound in rat. *Journal of the Korean Society of Food Science and Nutrition*, 43(4), 516-521.

Kim, J. S., Kwon, Y. S., Sa, Y. J., & Kim, M. J. (2011). Isolation and identification of sea buckthorn (*Hippophae rhamnoides*) phenolics with antioxidant activity and α -glucosidase inhibitory effect. *Journal of Agricultural and Food Chemistry*, 59(1), 138-144.

Ko, H. M., Eom, T. K., Song, S. K., Jo, G. Y., & Kim, J. S. (2017). Tyrosinase and α -glucosidase inhibition activities and antioxidant effects of extracts from different parts of *Hypochaeris radicata*. *Korea Journal of Medicinal Crop Science*, 25(3), 139-145.

Lee, H. J., Lim, G. N., Park, M. A., & Park, S. N. (2011). Antibacterial and antioxidative activity of *Lespedeza cuneata* G. Don extracts. *Korean Journal of Microbiology and Biotechnology*, 39(1), 63-69.

Numata, A., Hokimoto, K., & Yamaguchi, H. (1980). C-glycosylflavones in *Lespedeza cuneata*. *Chemical and Pharmaceutical Bulletin*, 28(3), 964-965.

Rhie, G. E., Shin, M. H., Seo, J. Y., Choi, W. W., Cho, K. H., Kim, K. H., Park, K. C., Eun, H. C., & Chung, J. H. (2001). Aging- and photoaging-dependent changes of enzymic and nonenzymic antioxidants in the epidermis and dermis of human skin *in vivo*. *Journal of Investigative Dermatology*, 117(5), 1212-1217.

Senguttuvan, J., Paulsamy, S., & Karthika, K. (2014). Phytochemical analysis and evaluation of leaf and root parts of the medicinal herb, *Hypochaeris radicata* L. for *in vitro* antioxidant activities. *Asian Pacific Journal of Tropical Biomedicine*, 4(Suppl 1), S359-S367.

Stich, H. F. (1991). The beneficial and hazardous effects of simple phenolic compounds. *Mutation Research/*

- Genetic Toxicology, 259(3-4), 307-324.
- Tsuji, N., Moriwaki, S., Suzuki, Y., Takema, Y., & Imokawa, G. (2001). The role of elastases secreted by fibroblasts in wrinkle formation: implication through selective inhibition of elastase activity. *Photochemistry and Photobiology*, 74(2), 283-290.
- Wojdylo, A., Oszmiański, J., & Czemerys, R. (2007). Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chemistry*, 105(3), 940-949.
- Zheng, W., & Wang, S. Y. (2001). Antioxidant activity and phenolic compounds in selected herbs. *Journal of Agricultural and Food Chemistry*, 49(11), 5165-5170.